

QHI Theory, II: Dilogarithmic and Quantum Hyperbolic Invariants of 3-Manifolds with $PSL(2, \mathbb{C})$ -Characters

[Abridged Version]

Stéphane Baseilhac and Riccardo Benedetti

Dipartimento di Matematica, Università di Pisa, Via F. Buonarroti, 2, I-56127 PISA. Emails: baseilha@mail.dm.unipi.it, benedett@dm.unipi.it.

The first author benefits from a Marie Curie Fellowship, contract No MCFI-2001-00605.

Abstract

We give parallel constructions of an invariant $R(W, \rho)$, based on the classical Rogers dilogarithm, and of quantum hyperbolic invariants (QHI), based on the Faddeev-Kashaev quantum dilogarithms, for flat $PSL(2, \mathbb{C})$ -bundles ρ over closed oriented 3-manifolds W . All these invariants are explicitly computed as a sum or state sums over the same hyperbolic ideal tetrahedra of the idealization of any fixed simplicial 1-cocycle description of (W, ρ) of a special kind, called a $\mathcal{D}$ -triangulation. $R(W, \rho)$ recovers the volume and the Chern-Simons invariant of ρ , and we conjecture that it determines the semi-classical limit of the QHI.

Keywords: state sum invariants, Cheeger-Chern-Simons invariants, dilogarithms, scissors congruences, volume conjecture.

1 Introduction

This text is a summary of our recent results. We give here with full details the complete apparatus of definitions, constructions and statements, with comments on the key points explaining the logic of our work. The proofs will be given in a forthcoming expanded version of this paper. Our aim here is to allow the interested reader to have, as much as possible, a quick access to these results. Concerning the (omitted) proofs, they are basically of two kinds: either verifications via careful computations with certain special functions, or geometric manipulations of (suitably decorated) triangulations of 3-manifolds. For the second ones, as well as for many properties of the special functions, one has, in fact, to repeat with mild modifications the arguments that are explained in full details in {Sections 1-6 + the Appendix} of [2]. The present paper also accomplishes the ‘ideology’ underlying Sections 7-9 of [2], which are, by the way, completely surpassed. The results of the present paper, limited to characters with values in a Borel subgroup of $PSL(2, \mathbb{C})$, had been announced in [4].

In Section 2, for any compact closed oriented 3-manifold W equipped with a character ρ with values in $PSL(2, \mathbb{C})$, one introduces special descriptions of (W, ρ) via simplicial $PSL(2, \mathbb{C})$ -valued 1-cocycles, called $\mathcal{D}$ -triangulations. A simple and geometrically meaningful procedure of *idealization* converts every $\mathcal{D}$ -triangulation $\mathcal{T}$ into an $\mathcal{I}$ -triangulation $\mathcal{T}_{\mathcal{I}}$ made by hyperbolic ideal tetrahedra with ordered vertices.

In Section 3, one constructs a suitably uniformized version R of the classical *Rogers dilogarithm* L . We interpret R as a function of ideal tetrahedra with ordered vertices, enriched with suitable $\mathbb{Z}$ -valued decorations of the edges that induce *log-branches* on the triples of moduli. It behaves well up to tetrahedral symmetries, and it verifies nice functional five term identities which lift arbitrary $2 \leftrightarrow 3$ moves on ideal triangulations. This section is largely inspired by W. Neumann's papers [19, 20].

In Section 4, one states that the signed sum $R(\mathcal{T}_{\mathcal{I}}, f)$ of the R -values of the tetrahedra of any fixed idealized triangulation $\mathcal{T}_{\mathcal{I}}$ (suitably enriched with a global *flattening* f , which induces a coherent log-branch system) well defines an invariant $R(W, \rho) \in \mathbb{C}/(\pi^2/6)\mathbb{Z}$, called the *dilogarithmic invariant*. This refines the $\mathbb{C}/\pi^2\mathbb{Q}$ -valued Dupont-Sah's description of the second Cheeger-Chern-Simons class for flat $SL(2, \mathbb{C})$ -bundles. In particular, $R(W, \rho)$ recovers the volume and the Chern-Simons invariant of the character ρ . We stress that the proof of invariance does not use any sophisticated group-cohomological argument: one shows directly that any two arbitrary (enriched) $\mathcal{I}$ -triangulations for (W, ρ) can be connected by so-called (enriched) $\mathcal{I}$ -transits. These transits are supported by usual elementary moves on bare triangulations, and dominated by $\mathcal{D}$ -transits between $\mathcal{D}$ -triangulations.

The dilogarithmic invariant emerged as a natural 'classical' counterpart from the effort in understanding the structure of the *quantum hyperbolic invariants* (QHI), that we had constructed in {Sections 1-6 + the Appendix} of [2] for triples (W, L, ρ) , where L is a *non-empty* link in W and ρ is only a B -valued character (B is the Borel subgroup of $SL(2, \mathbb{C})$ of upper triangular matrices). The main ingredient of the QHI are the Faddeev-Kashaev (non symmetric) *quantum dilogarithms*. In Section 5, one revisits the B -QHI construction in the spirit of the dilogarithmic invariant's one, and one realizes that:

(1) For any $PSL(2, \mathbb{C})$ -character ρ , by using the quantum dilogarithms one can define state sums supported by the idealization $\mathcal{T}_{\mathcal{I}}$ of any $\mathcal{D}$ -triangulation $\mathcal{T}$ for (W, ρ) . These state sums do not yet define an invariant because they do not behave well up to tetrahedral symmetries, and they verify only some special instances of five term identities.

(2) A specific procedure of *local* symmetrization of the quantum dilogarithms leads to fix an arbitrary non-empty link L in W in order to fix one coherent globalization. This is supported by any $\mathcal{D}$ -triangulation $\mathcal{T}$ for (W, ρ) in which the link L is realized as a Hamiltonian subcomplex H . The symmetrization of the quantum dilogarithms is governed by any fixed *integral charge* c relative to H . A charge is formally similar to the flattenings used for the dilogarithmic invariant, the main difference being that it is moduli-independent. It has a global structure which depends on the link-fixing, and eventually encodes the link itself. All this gives the notion of $\mathcal{D}$ -triangulation $(\mathcal{T}, H, c)$ for the triple (W, L, ρ) , and any such triangulation supports suitably modified state sums $H_N(\mathcal{T}_{\mathcal{I}}, c)$ indexed by the odd integers $N > 1$.

(3) The modified state sums satisfy all instances of five term identities, so they eventually well define (up to a sign and a N -th root of unity multiplicative factor) the QHI $H_N(W, L, \rho)$ for pairs (W, L) equipped with arbitrary $PSL(2, \mathbb{C})$ -characters.

Every state sum $H_N(\mathcal{T}_{\mathcal{I}}, c)$ looks formally very like $\exp(R(\mathcal{T}_{\mathcal{I}}, f))$, with the charge c playing the role of the flattening f . The presence of the link in the construction of

$H_N(W, L, \rho)$ is entirely a consequence of the adopted specific symmetrization procedure of the quantum dilogarithms. In order to get the general QHI for arbitrary $PSL(2, \mathbb{C})$ -characters, no further ‘quantum algebra’ than for the B -QHI is necessary. In fact the old B -state sums slightly differ from $H_N(\mathcal{T}, c)$ by a scalar factor which is expressed in terms of $(\mathcal{T}, c)$, not only of $(\mathcal{T}, c)$.

In Section 6 one places the dilogarithmic invariant and the QHI in the framework of the theory of *scissors congruence classes*.

One expects that the dilogarithmic invariant plays a main role to express the (dominant term of the) asymptotic expansion of the QHI when $N \rightarrow \infty$. This *Volume Conjecture* for $PSL(2, \mathbb{C})$ -QHI is discussed in Section 8 where it is also supported by some results about the actual asymptotic behaviour of the quantum dilogarithms.

In Section 9 one considers the extension of the dilogarithmic invariant and of the QHI to *cusped* hyperbolic 3-manifolds. This leads us to discuss the relationship between the corresponding Volume Conjecture and a current Volume Conjecture for the colored Jones invariant $J_N(L)$ of hyperbolic knots L in S^3 .

Finally we stress that our technic of ‘straightening’ (generic) 1-cocycles, which applies to $(PSL(2, \mathbb{C}), \partial \mathbb{H}^3)$ in the present paper and produces the idealization, looks very general. It can be considered for other pairs (G, X) , where G is a Lie group acting transitively on X and there is a suitable notion of *straight* tetrahedra with vertices in X . Cases of most interest are $(SU(2), S^3)$, $(SO(3), S^3)$, $(PSL(2, \mathbb{C}), \mathbb{H}^3)$. Moreover it should be interesting to extend it to a treatment of ‘truncated tetrahedra’ (hence manifolds with boundary). By using other ‘potential functions’ one could obtain the analogue of the dilogarithmic invariant for W endowed with G -valued characters. This could help to understand (and possibly refine) other state sum quantum-invariants, such as the Turaev-Viro one, in terms of scissors congruence classes (and, in particular, give a geometric interpretation to their asymptotic behaviour). We plan to study these generalizations in future works.

2 $\mathcal{D}$ -triangulations and the idealization

Let W be a compact, closed, oriented 3-manifold, and ρ be a flat principal bundle over W with structural group $PSL(2, \mathbb{C})$. We consider ρ up to isomorphisms of flat bundles; equivalently, it can be identified with a conjugacy class of representations in $PSL(2, \mathbb{C})$ of the fundamental group of W , i.e. with a $PSL(2, \mathbb{C})$ -character of W . The pairs (W, ρ) are considered up to oriented homeomorphisms. By using the *hauptvermutung*, depending on the context, we will freely assume that W is endowed with a (necessarily unique) PL or smooth structure, and use differentiable or PL homeomorphisms.

Given any singular triangulation T of W (i.e. a triangulation where tetrahedra may have self and multiple adjacencies), and any system of orientations of the edges of T , ρ can be represented by $PSL(2, \mathbb{C})$ -valued simplicial 1-cocycles, up to coboundaries of 0-cochains. For our purposes, we need to specialize the kind of triangulations, orientations and 1-cocycles.

Any T as above can be considered as a finite family $\{\Delta_i\}$ of *oriented abstract* tetrahedra, each being endowed with the standard triangulation with 4 vertices and the orientation induced by the one of W , together with a system of identifications of pairs of distinct (abstract) 2-faces. We will often distinguish between vertices, edges, 2-faces *in* T , that is after the identifications, and *abstract* edges,..., that is of the abstract Δ_i ’s. We view each Δ_i as positively embedded as a straight tetrahedron in $\mathbb{R}^3$ endowed with the orientation specified by the standard basis. Later we shall use

the hyperbolic space $\mathbb{H}^3$. It is also oriented by stipulating that, in the disk model, it is oriented as an open set of $\mathbb{R}^3$. The natural boundary $\partial\mathbb{H}^3 = \mathbb{CP}^1 = \mathbb{C} \cup \{\infty\}$ of $\mathbb{H}^3$ is oriented by its complex structure. The action of $PSL(2, \mathbb{C})$ on $\mathbb{H}^3$ and $\mathbb{CP}^1$ is the one of orientation preserving isometries of $\mathbb{H}^3$.

Definition 2.1 A $\mathcal{D}$ -triangulation for the pair (W, ρ) consists of a triple $\mathcal{T} = (T, b, z)$ where:

- (i) T is a *quasi-regular* triangulation of W , that is every edge of T has two distinct vertices as endpoints;
- (ii) b is a *branching* of T , that is a system of orientations of the edges such that the one induced on each abstract Δ_i is associated to a total ordering v_0, v_1, v_2, v_3 of its (abstract) vertices: each edge is oriented by the arrow emanating from the smallest endpoint;
- (iii) z is a 1-cocycle on (T, b) representing ρ such that (T, b, z) is *idealizable* (see Def. 2.3).

Remark 2.2 Quasi-regular (even regular indeed) triangulations of W do exist. A total ordering of the vertices of a quasi-regular triangulation T clearly induces a branching. *In the present paper we will only consider these special branchings associated to total orderings of the vertices.* This simplifies certain proofs, but all the results eventually hold true also for arbitrary branchings. For more information about branchings, see [2] and [6].

Given a branching b on a oriented tetrahedron Δ (realized in $\mathbb{R}^3$ as above), one can define an orientation of any of its simplices, not only of the edges. Denote by $E(\Delta)$ the set of b -oriented edges of Δ , and by e' the edge opposite to e . We put $e_0 = [v_0, v_1]$, $e_1 = [v_1, v_2]$ and $e_2 = [v_0, v_2] = -[v_2, v_0]$. The ordered triple of edges

$$(e_0 = [v_0, v_1], e_2 = [v_0, v_2], e'_1 = [v_0, v_3])$$

departing from v_0 defines a b -orientation of Δ . We say that (Δ, b) is positive if its b -orientation agrees with the one of $\mathbb{R}^3$, and negative otherwise; we indicate it by a sign $*$ = $*_b = \pm 1$. The 2-faces of Δ can be named by their opposite vertices. We orient them by working as above on the boundary of each 2-face f : there is a b -ordering of the vertices of f , and an orientation of f which induces on ∂f the prevailing orientation among the three b -oriented edges. If z is $PSL(2, \mathbb{C})$ -valued 1-cocycle on (Δ, b) , we write $z_j = z(e_j)$ and $z'_j = z(e'_j)$. Then, one reads the cocycle condition on the 2-face opposite to v_3 as $z_0 z_1 z_2^{-1} = 1$. These considerations apply to each abstract tetrahedron of any branched triangulation (T, b) of W and to (the restrictions of) any $PSL(2, \mathbb{C})$ -valued 1-cocycle z on (T, b) .

Definition 2.3 Let (Δ, b, z) be a branched tetrahedron endowed with a $PSL(2, \mathbb{C})$ -valued 1-cocycle z . It is *idealizable* iff

$$u_0 = 0, \quad u_1 = z_0(0), \quad u_2 = z_0 z_1(0), \quad u_3 = z_0 z_1 z'_0(0)$$

are 4 distinct points in $\mathbb{C} \subset \mathbb{CP}^1 = \partial\bar{\mathbb{H}}^3$ which span an non degenerate hyperbolic ideal tetrahedron (with ordered vertices). A triangulation (T, b, z) is *idealizable* iff all its tetrahedra (Δ_i, b_i, z_i) are idealizable.

If (Δ, b, z) is idealizable, for all $j = 0, 1, 2$ one can associate to e_j and e'_j the same *cross-ratio* modulus $w_j \in \mathbb{C} \setminus \{0, 1\}$ of the hyperbolic ideal tetrahedron defined by (u_0, u_1, u_2, u_3) . By using the cyclic ordering of the edges induced by b , one has

$$w_{j+1} = 1/(1 - w_j)$$

and

$$w_0 = (u_2 - u_1)u_3/u_2(u_3 - u_1) := -p_1/p_2 .$$

Similarly we shall write

$$w_j = -p_{j+1}/p_{j+2} .$$

Set $w = (w_0, w_1, w_2)$ and call it a *modular triple*. As the ideal tetrahedron is non-degenerate, the imaginary parts of the w_j 's are not equal to zero, and share the same sign $*_w = \pm 1$.

Definition 2.4 We call (Δ, b, w) the *idealization* of the idealizable (Δ, b, z) . For any $\mathcal{D}$ -triangulation $\mathcal{T} = (T, b, z)$ of (W, ρ) , its *idealization* $\mathcal{T}_{\mathcal{I}} = (T, b, w)$ is given by the family $\{(\Delta_i, b_i, w_i)\}$ of idealizations of the (Δ_i, b_i, z_i) 's. We say that $\mathcal{T}_{\mathcal{I}}$ is an $\mathcal{I}$ -triangulation.

Remarks 2.5 (1) We have inglobed the ‘non-degenerate’ assumption into the notion of ‘idealizable’. This simplifies the exposition and also certain proofs; however, it is not necessary for the validity of the results of the paper.

(2) If z_j acts on $\mathbb{CP}^1$ as $(a_j x + b_j)/(c_j x + d_j)$, then $z_j(0) = b_j/d_j$. Thus it is immediate to formulate the ‘idealizability’ condition in terms of a simple system of real algebraic inequalities on the entries of the z_j 's.

(3) In [2] we have used so-called *full* B -valued 1-cocycles z to construct the B -QHI. This means that for any edge e the upper-diagonal entry $x(e)$ of $z(e)$ is non-zero. It is easy to verify that a B -cocycle is full iff it is idealizable (forgetting the non-degenerate assumption). In [4] we proposed an idealization of full B -cocycles, which was indeed a specialization of the present general procedure. The idealization of a $\mathcal{D}$ -tetrahedron with a full B -cocycle is simply given by $w_j = -q_{j+1}/q_{j+2}$, where $q_j = x(e_j)x(e'_j)$ for $j = 0, 1$, and $q_2 = -x(e_2)x(e'_2)$.

(4) It follows from the cocycle condition that $p_0 + p_1 + p_2 = 0$ (also that $q_0 + q_1 + q_2 = 0$ for full B -cocycles).

The following lemma is immediate:

Lemma 2.6 *For any $PSL(2, \mathbb{C})$ -character ρ , any quasi-regular branched triangulation (T, b) of W can be completed to a $\mathcal{D}$ -triangulation (T, b, z) of the pair (W, ρ) .*

In fact, given any 1-cocycle, one can perturb it by the coboundaries of generic 0-cochains which are injective on the vertices of T , and get an idealizable one.

Tetrahedral symmetries. The following simple lemma states the good behaviour of the idealization with respect to a change of branching (i.e. the ‘tetrahedral symmetries’).

Lemma 2.7 *Denote by S_4 the permutation group on four elements. A permutation $p \in S_4$ of the vertices of an idealizable tetrahedron (Δ, b, z) gives another idealizable tetrahedron (Δ, b', z') . For every edge e , $z'(e) = z(e)^{-1}$ iff the orientations of e for b and b' are opposite; otherwise $z'(e) = z(e)$. The permutation turns the idealization (Δ, b, w) into (Δ, b', w') , where, for each edge e , $w'(e) = w(e)^{\epsilon(p)}$, and $\epsilon(p)$ is the signature of p .*

Consider for instance the transposition $(0, 1)$. It turns the set of 4 points

$$0, z_0(0), z_0 z_1(0), z_0 z_1 z'_0(0)$$

into

$$0, (z_0)^{-1}(0), z_1(0), z_1 z'_0(0) .$$

By applying on the second set of points the hyperbolic isometry z_0 one gets the initial set after the transposition of its first two members, and similarly for other permutations. The lemma follows immediately, due to the behaviour of the cross-ratio up to vertex permutation.

Remark 2.8 As the signature of a permutation also changes the sign $*_b$ of the branching, the above lemma is coherent with the usual symmetry relations holding in the classical (pre)-Bloch group: $[x] = [1/(1-x)]$ and $[x] = -[1/x]$ (see e.g. [10], [19]).

Hyperbolic edge compatibility. We are now concerned with an important global property of the idealized triangulations $\mathcal{T}_{\mathcal{I}}$. Before to state it, let us stress that when dealing with modular triples one has to be careful with the orientations. Recall that every $\mathcal{I}$ -tetrahedron (Δ, b, w) is oriented by definition; in the case of an $\mathcal{I}$ -triangulation this is given by the orientation of W . There is also the b -orientation encoded by the sign $* = *_b$. The idealization ‘physically’ realizes the vertices of Δ on $\partial\mathbb{H}^3$, with the ordering induced by b . So the b -orientation may or may not agree with the orientation of the spanned ideal tetrahedron induced by the fixed orientation of $\mathbb{H}^3$, which is encoded by the sign $*_w$ of the modular triple.

Definition 2.9 We say that (Δ, b, w) is *geometric* iff $*_b = *_w$.

Given any $\mathcal{I}$ -triangulation $\mathcal{T}_{\mathcal{I}} = (T, b, w)$, the contribution of each (Δ_i, b_i, w_i) to any computation with the moduli is given by the $w(e)^*$ ’s, where e is any edge in Δ_i and $* = *_b$. The next Lemma 2.10 is a first concretization of this fact (see also the notion of $\mathcal{I}$ -transit below).

Let $\mathcal{T}_{\mathcal{I}} = (T, b, w)$ be as above. Denote by $E(T)$ the set of edges of T , by $E_{\Delta}(T)$ the whole set of edges of the associated abstract tetrahedra $\{\Delta_i\}$, and by $\epsilon : E_{\Delta}(T) \longrightarrow E(T)$ the natural identification map.

Lemma 2.10 For any edge $e \in E(T)$, we have $\prod_{a \in \epsilon^{-1}(e)} w(a)^* = 1$, where $* = \pm 1$ according to the b -orientation of the tetrahedron Δ_i that contains a .

This means that the $w(a)^*$ ’s around each e verify the usual compatibility condition, necessary when one tries to construct hyperbolic 3-manifolds by glueing ideal tetrahedra. A key point in the proof is that around every edge e of T one may only meet an even number of tetrahedra such that the b -orientations of the two faces containing e are opposite. This follows from the fact that W is orientable.

Remark 2.11 Every idealized triangulation $\mathcal{T}_{\mathcal{I}}$ for (W, ρ) necessarily includes some *non-geometric* ideal tetrahedra. In fact if all the tetrahedra were geometric, the compatibility condition in Lemma 2.10 would imply that the sum of the arguments is exactly equal to 2π around all the edges of T , not only an even multiple of 2π (see Lemma E.6.1. of [5]). So, one should have a genuine hyperbolic manifold structure on W out of the vertices of T , with W being triangulated by embedded geodesic ideal tetrahedra. Using tetrahedra truncated by suitable horospheres, one deduces that the spherical link surrounding each vertex of T would inherit a $(\mathbb{C}^*, \mathbb{C})$ -structure, which is impossible.

$\mathcal{D}$ - and $\mathcal{I}$ -transits. It is well-known that given any two (singular) triangulations T_0 and T_1 of W there exists a finite sequence $T_0 \rightarrow \dots \rightarrow T_1$ of so-called $2 \leftrightarrow 3$ and *bubble moves* that turns T_0 into T_1 . For instance, it is a consequence of the duality between ideal triangulations and standard spines of $T \setminus T^0$ (removing the vertices), and the calculus for standard spines due to Matveev [17] and Piergallini




Figure 1: the moves between singular triangulations.

[22]. It is sometimes convenient, for technical reasons (for instance when dealing with arbitrary branchings as in [2]), to consider a further so-called $0 \leftrightarrow 2$ move. See Fig. 1.

Next we consider moves on $\mathcal{D}$ -triangulations $\mathcal{T} = (T, b, z)$ and $\mathcal{I}$ -triangulations $\mathcal{T}_{\mathcal{I}} = (T, b, w)$ for the pair (W, ρ) , called $\mathcal{D}$ -transits and $\mathcal{I}$ -transits respectively. They are supported by the above bare triangulation moves, but they also include the transits of the respective extra-structures.

First of all one requires that the condition to be quasi-regular is preserved by the moves. We stress that this is not an automatic fact, on the contrary this leads to one main technical complication in the proofs.

A move $T_0 \leftrightarrow T_1$ between branched triangulations (T_0, b_0) and (T_1, b_1) defines a *branching transit* $(T_0, b_0) \leftrightarrow (T_1, b_1)$ if b_0 and b_1 agree on the common edges. As stipulated above, we shall only consider branched quasi-regular triangulations, where the branchings are defined by total orderings of the vertices. In such a case any $2 \leftrightarrow 3$ or $0 \leftrightarrow 2$ move that preserves the quasi-regularity of the triangulations can be completed in a unique way to a branching transit, but a bubble move may be completed in different ways, each of them being a possible transit.

Let $(T_0, b_0), (T_1, b_1)$ be as above and $z_k \in Z^1(T_k; PSL(2, \mathbb{C}))$, $k = 0, 1$. We have a (resp. *idealizable*) *cocycle transit* $(T_0, b_0, z_0) \leftrightarrow (T_1, b_1, z_1)$ if z_0 and z_1 agree on the common edges (resp. and both are idealizable 1-cocycles). Note that for $2 \rightarrow 3$ and $0 \rightarrow 2$ moves, given z_k there is only one (resp. at most one) z_{k+1} with this property. We stress that in some special cases a $2 \rightarrow 3$ transit of an idealizable cocycle can actually not preserve the idealizability, but generically this does not hold. For positive bubble moves there is always an infinite set of possible (idealizable) cocycle transits.

We say that $(T_0, b_0, z_0) \leftrightarrow (T_1, b_1, z_1)$ as above is a $\mathcal{D}$ -transit when both z_0 and z_1 are idealizable.

Let us now consider the transit for the idealized triangulations. Consider the convex hull of five distinct points $u_0, u_1, u_2, u_3, u_4 \in \partial\mathbb{H}^3$, with the two possible triangulations Q_0, Q_1 made of the oriented hyperbolic ideal tetrahedra Δ^i obtained by omitting u_i . An edge e of $Q_i \cap Q_{i+1}$ belongs to one tetrahedron of Q_i iff it belongs to two tetrahedra of Q_{i+1} . Then, the modulus of e in Q_i is the product of the two moduli of e in Q_{i+1} . Also, the product of the moduli on the central edge of Q_1 is equal to 1.

Let $T \rightarrow T'$ be a $2 \rightarrow 3$ move. Consider the two (resp. three) abstract tetrahedra of T (resp. T') involved in the move. They determine subsets $\tilde{E}(T)$ of $E_\Delta(T)$ and $\tilde{E}(T')$ of $E_\Delta(T')$. Denote their complementary sets by $\hat{E}(T)$ and $\hat{E}(T')$. Clearly one can identify $\hat{E}(T)$ and $\hat{E}(T')$. Using the above configurations Q_0 and Q_1 , and recalling the considerations made before Lemma 2.10, we are led to the following definition:




Figure 2: a $2 \leftrightarrow 3$ ideal transit.

Definition 2.12 One has a $2 \rightarrow 3$ $\mathcal{I}$ -transit $(T, b, w) \rightarrow (T', b', w')$ of $\mathcal{I}$ -triangulations for a pair (W, ρ) if:

- 1) w and w' agree on $\hat{E}(T) = \hat{E}(T')$;
- 2) for each common edge $e \in \epsilon_T(\tilde{E}(T)) \cap \epsilon_{T'}(\tilde{E}(T'))$ one has

$$\prod_{a \in \epsilon_T^{-1}(e)} w(a)^* = \prod_{a' \in \epsilon_{T'}^{-1}(e)} w'(a')^* , \quad (1)$$

where $*$ = ± 1 according to the b -orientation of the tetrahedron that contains a (resp. a'). One has a $0 \rightarrow 2$ (resp. *bubble*) $\mathcal{I}$ -transit if the above first condition is satisfied, and one replaces the second by:

- 2') for each edge $e \in \epsilon_{T'}(\tilde{E}(T'))$ one has

$$\prod_{a' \in \epsilon_{T'}^{-1}(e)} w'(a')^* = 1 . \quad (2)$$

$\mathcal{I}$ -transits for negative $3 \rightarrow 2$ moves are defined in exactly the same way, and for negative $2 \rightarrow 0$ and bubble moves w' is defined by simply forgetting the moduli of

the two disappearing tetrahedra. The condition (1) above, implies that the product of the $w'(a')^*$'s around the new edge is equal to 1. A $2 \leftrightarrow 3$ $\mathcal{I}$ -transit is shown in Fig. 2; we only indicate the first component of each modular triple. In general, the relations (1) may imply that w or w' equals 0 or 1 on some edges. In that case, the $2 \leftrightarrow 3$ $\mathcal{I}$ -transit fails. In particular, in Fig. 2 we assume that $x \neq y$.

Note that for $2 \leftrightarrow 3$ $\mathcal{I}$ -transits w' is uniquely determined by w , whereas for $0 \rightarrow 2$ and bubble $\mathcal{I}$ -transits there is one degree of freedom in choosing w' . Also, condition (2) simply means that such transits give *the same* modular triples to the two new tetrahedra, for their b -orientations are opposite.

Remark 2.13 Once it is expressed in terms of the involved moduli w_0 's, any $2 \leftrightarrow 3$ $\mathcal{I}$ -transit dominates an instance of the *five term identities* which enter in the construction of the classical (pre)-Bloch group. For example, the transit in Fig. 2 corresponds to the relation

$$[x] - [y] + [y/x] - [(1 - x^{-1})/(1 - y^{-1})] + [(1 - x)/(1 - y)] = 0 .$$

The next proposition states the remarkable fact that $\mathcal{D}$ -transits and $\mathcal{I}$ -transits together with the idealization make commutative diagrams, that is the $\mathcal{D}$ -transits *dominate* the $\mathcal{I}$ -transits.

Proposition 2.14 *Let $\mathfrak{d}$ be any $\mathcal{D}$ -transit of $\mathcal{D}$ -triangulations and $\mathfrak{i}$ be any $\mathcal{I}$ -transit of $\mathcal{I}$ -triangulations for a pair (W, ρ) . Denote by $\mathcal{I}$ the idealization map $\mathcal{T} \rightarrow \mathcal{T}_{\mathcal{I}}$. Then for every $\mathfrak{d}$ there exists $\mathfrak{i}$ (resp. for every $\mathfrak{i}$ there exists $\mathfrak{d}$) such that $\mathfrak{i} \circ \mathcal{I} = \mathcal{I} \circ \mathfrak{d}$.*

For $2 \leftrightarrow 3$ transits there is also an uniqueness statement. The proof is not hard. By using the tetrahedral symmetries of Lemma 2.7, it is enough to show the proposition for one branching transit configuration (for instance the one of Fig. 2). The verification follows almost immediately from the definition of the idealization, as for Lemma 2.7. Note that the possible failures of $2 \rightarrow 3$ transits of idealizable cocycles that we mentionned above correspond to the failures of $2 \rightarrow 3$ $\mathcal{I}$ -transits (for instance when $x = y$ in Fig. 2).

3 Classical dilogarithms

3.1 Rogers dilogarithm

Denote by $\log$ the standard branch of the logarithm, with arguments in $] - \pi, \pi]$. Put $\mathfrak{D} = \mathbb{C} \setminus \{(-\infty; 0) \cup (1; +\infty)\}$. The *Rogers dilogarithm* is the complex analytic function defined over $\mathfrak{D}$ by

$$L(x) = -\frac{\pi^2}{6} - \frac{1}{2} \int_0^x \left(\frac{\log(t)}{1-t} + \frac{\log(1-t)}{t} \right) dt , \quad (3)$$

where we integrate first along the path $[0; 1/2]$ on the real axis and then along any path in $\mathfrak{D}$ from $1/2$ to x . Here we add $-\pi^2/6$ so that $L(1) = 0$. For $|x| < 1$, one may also write L as

$$L(x) = -\frac{\pi^2}{6} + \frac{1}{2} \log(x) \log(1-x) + \sum_{n=1}^{\infty} \frac{x^n}{n^2} .$$

The sum in the right-hand side is the power series expansion in the open unit disk of the *Euler dilogarithm* Li_2 , defined over $\mathbb{C} \setminus (1; +\infty)$ by

$$\text{Li}_2(x) = - \int_0^x \frac{\log(t)}{1-t} dt .$$

For a detailed study of the dilogarithm functions and their relatives, see [16] or the review [24]. Below we will also call ‘Rogers dilogarithm’ the multi-valued function obtained from L via analytic continuation. The function L is related to the *Bloch-Wigner dilogarithm*

$$D_2(x) = \text{Im}(\text{Li}_2(x)) + \arg(1-x) \log|x| ,$$

which is obtained by adding to $\text{Im}(\text{Li}_2(x))$ the needed correction term to compensate its jump along the branch cut $(1; +\infty)$. The function $D_2(x)$ is a real analytic continuation of $\text{Im}(\text{Li}_2(x))$ on $\mathbb{C} \setminus \{0, 1\}$, and it is continuous (but not differentiable) at 0 and 1. It gives the volume of oriented hyperbolic tetrahedra by the formula

$$\text{Vol}(\Delta, b, w) = D_2(w_0) , \quad (4)$$

where, with the notations of Section 2, w denotes the modular triple with respect to a positive branching b . In particular, we have the 6-fold symmetries

$$D_2(w_0) = D_2(w_1) = D_2(w_2) = -D_2(w_0^{-1}) = -D_2(w_1^{-1}) = -D_2(w_2^{-1}) . \quad (5)$$

Moreover, if we apply the formula (4) to Fig. 2 we get the five term functional relation

$$D_2(y) + D_2\left(\frac{1-x^{-1}}{1-y^{-1}}\right) = D_2(x) + D_2(y/x) + D_2\left(\frac{1-x}{1-y}\right) \quad (6)$$

when $x \neq y$. Finally, all the other five term relations obtained by changing the branching in Fig. 2 also hold true, due to (5).

One would like to think of the Rogers dilogarithm L as the natural complex analytic analogue of $D_2(x)$. But L verifies similar five term relations only by putting restrictions on the variables. For instance, the analog of (6) is the so-called Schaeffer’s identity

$$L(x) - L(y) + L(y/x) - L\left(\frac{1-x^{-1}}{1-y^{-1}}\right) + L\left(\frac{1-x}{1-y}\right) = 0 \quad (7)$$

which for real x, y holds only when $0 < y < x < 1$. In fact, this identity characterizes the Rogers dilogarithm: if $f(0; 1) \rightarrow \mathbb{R}$ is a 3 times differentiable function satisfying (7) for all $0 < y < x < 1$, then $f(x) = kL(x)$ for a suitable constant k [23, Sect. 4], [9, App.]. By analytic continuation, the relation (7) holds true for complex parameters x, y , providing that the imaginary part of y is ≥ 0 , and x lies inside the triangle formed by 0, 1 and y . For such x, y , note that also all the other arguments of L in (7) have positive imaginary parts. With this restriction, this relation corresponds to one specific instance of $\mathcal{I}$ -transit (see Remark 2.13). This naively suggests the possibility to set

$$L(\Delta, b, w) = L(w_0) ,$$

and try to use it to build an invariant for (W, ρ) that should be computable by using any idealized triangulation $\mathcal{T}_L$. However, one is immediately faced to the following correlated difficulties:

- (1) (*Uniformization*) One has to deal with the different branches of L .
- (2) (*Symmetrization*) One realizes that $L(\Delta, b, w)$ only respects the tetrahedral symmetries up to some elementary functions (see below).
- (3) (*Completing the five term relations*) As remarked above, L satisfies certain five term relations which correspond only to some peculiar instances of $\mathcal{I}$ -transits, and with restrictions on the range of moduli.

In the next three subsections we outline how to solve these difficulties.

3.2 Uniformization

Let $\widehat{\mathbb{C}} = \widehat{\mathbb{C}}_{00} \cup \widehat{\mathbb{C}}_{01} \cup \widehat{\mathbb{C}}_{10} \cup \widehat{\mathbb{C}}_{11}$, where $\widehat{\mathbb{C}}_{\varepsilon\varepsilon'}$ ($\varepsilon, \varepsilon' = 0, 1$) is the Riemann surface of the function defined on $\mathfrak{D} = \mathbb{C} \setminus \{(-\infty; 0) \cup (1; +\infty)\}$ by

$$x \mapsto (\log(x) + \varepsilon i\pi, \log((1-x)^{-1}) + \varepsilon' i\pi) .$$

Thus $\widehat{\mathbb{C}}$ is the abelian cover of $\mathbb{C} \setminus \{0, 1\}$ obtained from $\mathfrak{D} \times \mathbb{Z}^2$ by the identifications

$$\begin{aligned} \{(-\infty; 0) + i0\} \times \{p\} \times \{q\} &\sim \{(-\infty; 0) - i0\} \times \{p+2\} \times \{q\} \\ \{(1; +\infty) + i0\} \times \{p\} \times \{q\} &\sim \{(1; +\infty) - i0\} \times \{p\} \times \{q+2\} , \end{aligned}$$

and the function

$$l(x; p, q) = (\log(x) + pi\pi, \log((1-x)^{-1}) + qi\pi) \quad (8)$$

is well-defined and analytic on $\widehat{\mathbb{C}}$. Here $(-\infty; 0) \pm i0$ comes from the upper/lower fold of $\mathfrak{D}$ with respect to $(-\infty; 0)$, and similarly for $(1; +\infty) \pm i0$. Following [19, 20], consider the following lift of the Rogers dilogarithm on $\widehat{\mathbb{C}}$:

$$R(x; p, q) = L(x) + \frac{i\pi}{2}(p \log(1-x) + q \log(x)) . \quad (9)$$

Lemma 3.1 *The above formula well defines an analytic map $R : \widehat{\mathbb{C}} \rightarrow \mathbb{C}/\pi^2\mathbb{Z}$.*

One can view R as a uniformization $\text{mod}(\pi^2)$ of L . We want to interpret R as a function of our $\mathcal{I}$ -tetrahedra (Δ, b, w) . In order to do that, it is natural to enrich the decoration by a $\mathbb{Z}$ -valued function f on the edges of Δ such that, for every edge, $f(e) = f(e')$. As for $w = (w_0, w_1, w_2)$, one can write $f = (f_0, f_1, f_2)$ with respect to b . Then we set

$$R(\Delta, b, w, f) = R(w_0; f_0, f_1) .$$

Next we indicate under which condition on f the function $R(\Delta, b, w, f)$ respects the tetrahedral symmetries.

3.3 Tetrahedral symmetries

Here is a crucial definition.

Definition 3.2 Let (Δ, b, w, f) and $f = (f_0, f_1, f_2)$ be as above. Set

$$l_j(b, w, f) = \log(w_j) + f_j i\pi ,$$

for $j = 1, 2, 3$. We say that (f_0, f_1, f_2) is a *flattening* of (Δ, b, w) if

$$l_0(b, w, f) + l_1(b, w, f) + l_2(b, w, f) = 0 .$$

In that case, we call $l_j(b, w, f)$ a *log-branch* of (Δ, b, w) for the edge e_j , and set $l(b, w, f) = (l_0(b, w, f), l_1(b, w, f), l_2(b, w, f))$ for the total log-branch associated to f .

Note that if f is a flattening of (Δ, b, w) , then it is a flattening of (Δ, b, u) for every modular triple u sufficiently close to w .

Let (Δ, b', w, f) be any enriched $\mathcal{I}$ -tetrahedron. By acting with a permutation $p \in S_4$ on the vertices of Δ , one passes from b' to a new branching b . Denote by (Δ, b, w, f) the new tetrahedron, where one still associates to every edge the same values $w(e)$ and $f(e)$ as before, but they are renamed according to the new ordering of the vertices given by b . Let $\epsilon(p)$ be the signature of p . Set $w^{\epsilon(p)} = (w_0^{\epsilon(p)}, w_1^{\epsilon(p)}, w_2^{\epsilon(p)})$ and $\epsilon(p)f = (\epsilon(p)f_0, \epsilon(p)f_1, \epsilon(p)f_2)$. We have

Lemma 3.3 *For any enriched $\mathcal{I}$ -tetrahedron (Δ, b, w, f) the identities*

$$R(\Delta, b', u, f) = \epsilon(p) R(\Delta, b, u^{\epsilon(p)}, \epsilon(p)f) \pmod{(\pi^2/6)\mathbb{Z}}$$

hold true for every permutation p and for every modular triple u sufficiently close to w if and only if f is a flattening of (Δ, b, w) .

3.4 Complete five term relations

We first define a notion of transit between flattened tetrahedra, and then use it to get all the required five term relations, without restrictions neither on the underlying branching transit configuration, nor on the range of the moduli.

Consider a $2 \rightarrow 3$ $\mathcal{I}$ -transit $(T, b, w) \rightarrow (T', b', w')$ as in Fig. 2. Give a flattening to each tetrahedron of the initial configuration, and denote by l the corresponding log-branch function on T . Recall the definition of the map ϵ_T before Lemma 2.10. The simple idea is just to formally take the log of the $\mathcal{I}$ -transit.

Definition 3.4 A map $f' : E_\Delta(T') \rightarrow \mathbb{Z}$ defines a $2 \rightarrow 3$ *log-branch transit* $(T, b, w, f) \rightarrow (T', b', w', f')$ if for each common edge $e \in T \cap T'$ one has the following relation between log-branches:

$$\sum_{a \in \epsilon_T^{-1}(e)} * l(a) = \sum_{a' \in \epsilon_{T'}^{-1}(e)} * l'(a') , \quad (10)$$

where $*$ = ± 1 according to the b -orientation of the tetrahedron that contains a (resp. a'). A map $f' : E_\Delta(T') \rightarrow \mathbb{Z}$ defines a $0 \rightarrow 2$ (resp. bubble) *log-branch transit* if for each edge $e \in T'$ one has

$$\sum_{a' \in \epsilon_{T'}^{-1}(e)} * l'(a') = 0 . \quad (11)$$

One easily verifies that log-branch transits actually define flattened tetrahedra, and that for a $2 \rightarrow 3$ log-branch transit the sum of values of l' about the new edge is always equal to zero. So log-branch transits for negative $3 \rightarrow 2$ moves are defined in exactly the same way, except that we also require that this last condition holds. For negative $2 \rightarrow 0$ and bubble moves the log-branch transits are defined by simply forgetting the log-branches of the two disappearing tetrahedra. The flattenings of a log-branch transit, associated to a given $\mathcal{I}$ -transit $(T, b, w) \rightarrow (T', b', w')$, actually define a log-branch transit for every $\mathcal{I}$ -transit $(T, b, u) \rightarrow (T', b', u')$ if u (resp. u') is a modular triple sufficiently close to w (resp. w').

Note that the relations (11) mean that the two new tetrahedra have *the same* log-branches, for their b -orientations are always opposite.

Proposition 3.5 *Let $(T, b, w, f) \rightarrow (T', b', w', f')$ be a log-branch transit. Then we have*

$$\sum_{\Delta \subset T} * R(\Delta, b, w, f) = \sum_{\Delta' \subset T'} * R(\Delta', b', w', f') \pmod{(\pi^2/6)\mathbb{Z}} , \quad (12)$$

where $$ = ± 1 according to the b -orientation of Δ (resp. Δ').*

Thanks to the tetrahedral symmetries, it is enough to prove the proposition for one branching transit configuration. For instance, for $2 \leftrightarrow 3$ transits one uses the one in Fig. 2, which underlies Schaeffer's identity (7). The key and delicate point consists in realizing that the log-branch transit condition determines a proper *analytic* subset $\mathfrak{G}$ of $\widehat{\mathbb{C}}^5$, on which the five term relation (12) corresponds to an *analytic* relation. Then one verifies that this relation holds true on a non empty open subset of $\mathfrak{G}$, as a consequence of Schaeffer's identity (7) over its complex domain of validity. Then the result follows by using the analytic continuation principle.

Remark 3.6 The function $R(x; p, q)$ with its five term relation that lifts Schaefer's identity was already considered by W. Neumann [19, 20]. A peculiarity of our treatment of the function $R(\Delta, b, w, f)$ is the preliminary discussion on the tetrahedral symmetries, from which the notion of log-branch emerges straightforwardly in a natural way (see Lemma 3.3). Also, when considering a $2 \leftrightarrow 3$ move configuration between hyperbolic ideal tetrahedra, we thus have to look at all possible branching transits, for $R(\Delta, b, w, f)$ explicitly depends on the branching b . Each branching transit leads to a specific instance of functional lifted five term relation for $R(\Delta, b, w, f)$, and there actually exist some which only hold mod $(\pi^2/6)\mathbb{Z}$.

4 The dilogarithmic invariant $R(W, \rho)$

Let $\mathcal{T} = (T, b, z)$ be a $\mathcal{D}$ -triangulation for (W, ρ) , with idealization $\mathcal{T}_{\mathcal{I}} = (T, b, w)$. In order to define the dilogarithmic invariant, one has to enrich $\mathcal{T}_{\mathcal{I}}$ with a system f of flattenings *stable* for log-branch transits.

Definition 4.1 We say that f is a *flattening* of $\mathcal{T}_{\mathcal{I}} = (T, b, w)$ if each f_i is a flattening of (Δ_i, b_i, w_i) and the following global conditions are verified:

- (1) The sum of b -signed log-branches around every edge of T is equal to 0.
- (2) Condition (1) implies that f verifies a similar one mod(2), so that it represents a class $[f] \in H^1(W, \mathbb{Z}/2\mathbb{Z})$. One requires that $[f] = 0$.

The first condition is formally the log of the compatibility condition in Lemma 2.10. The second ensures a nice *affine* structure on the set of these flattenings. A slight adaptation of a fundamental result due to W. Neumann [18, 20] implies the following:

Theorem 4.2 Any $\mathcal{I}$ -triangulation $\mathcal{T}_{\mathcal{I}} = (T, b, w)$ for (W, ρ) admits a flattening. Flattenings on $\mathcal{T}_{\mathcal{I}} = (T, b, w)$ make an affine space over a lattice. This lattice has an explicitly given basis made by one vector for the abstract star of each edge of T .

Finally one can state

Theorem 4.3 Let $\mathcal{T}_{\mathcal{I}}$ be any flattened $\mathcal{I}$ -triangulation for (W, ρ) . Set

$$R(\mathcal{T}_{\mathcal{I}}, f) = \sum_i *_{\mathbf{i}} R(\Delta_i, b_i, w_i, f_i) \in \mathbb{C}/(\pi^2/6)\mathbb{Z} ,$$

where $*_{\mathbf{i}} = \pm 1$ according to b_i . The value of $R(\mathcal{T}_{\mathcal{I}}, f)$ does not depend on the choice of $(\mathcal{T}_{\mathcal{I}}, f)$. Hence it defines an invariant $R(W, \rho) \in \mathbb{C}/(\pi^2/6)\mathbb{Z}$ called the dilogarithmic invariant of the pair (W, ρ) .

Proposition 4.4 We have $R(W, \rho) = \text{CS}(\rho) + i\text{Vol}(\rho) \bmod(\pi^2/6)\mathbb{Z}$, where CS and Vol are the Chern-Simons invariant and the volume of the character ρ .

This proposition shows that $R(W, \rho)$ refines the mod($\pi^2\mathbb{Q}$) dilogarithmic interpretation of the second Cheeger-Chern-Simons class of ρ , due to Dupont-Sah [8, 9]. This is in agreement (except for the mod($\pi^2/6$) $\mathbb{Z}$ precision) with the results stated in [19, 21] in the particular case when ρ is the holonomy of a genuine hyperbolic structure on W .

In [4] we had roughly announced the existence (limited to B -characters) of such dilogarithmic invariant defined mod($\pi^2/2$) $\mathbb{Z}$. A more careful analysis of the tetrahedral symmetries leads to the present mod($\pi^2/6$) $\mathbb{Z}$ formulation.

5 Quantum dilogarithms and QHI for (W, L, ρ)

In this section we revisit the construction of the B-QHI made in [2], having as model the construction of the dilogarithmic invariant. The result shall be a full generalization of the QHI for arbitrary $PSL(2, \mathbb{C})$ -characters, and a deeper understanding of the ultimate hyperbolic geometric nature of these invariants.

5.1 Quantum dilogarithms

Let $N > 1$ be a fixed odd positive integer, and put $\zeta = \exp(2i\pi/N)$. The *Faddeev-Kashaev's quantum dilogarithms* were originally derived as an explicit matrix realization of the associator in the cyclic representation theory of a Borel quantum subalgebra of $U_\zeta(sl(2, \mathbb{C}))$ (see [12, 13]). The matrix elements of the quantum dilogarithms are usually called $6j$ -symbols. This derivation is also presented in the Appendix of [2], and with full details in [1]. Here we forget this ‘quantum algebraic’ origin, and simply describe and interpret the special functions one has to deal with. For any complex number x with $|x| < 1$, consider the analytic function g defined by

$$g(x) := \prod_{j=1}^{N-1} (1 - x\zeta^j)^{j/N}$$

and set $h(x) := x^{-p}g(x)/g(1)$ when x is non-zero (one computes that $|g(1)| = N^{1/2}$). We shall still denote by g its analytic continuation to the complex plane with cuts from the points $x = \exp(i\epsilon)\zeta^k$, $k = 0, \dots, N-1$, $\epsilon \in \mathbb{R}$, to infinity. Hereafter we will implicitly assume that ϵ is such that the cuts are away from the points where g is evaluated.

Consider the curve $\Gamma = \{x^N + y^N = z^N\} \subset \mathbb{CP}^2$ (homogeneous coordinates), and the rational functions given for any $n \in \mathbb{N}$ by

$$\omega(x, y, z|n) = \prod_{j=1}^n \frac{(y/z)}{1 - (x/z)\zeta^j} \quad . \quad (13)$$

These functions are periodic in their integer argument, with period N . Denote by δ the N -periodic Kronecker symbol, i.e. $\delta(n) = 1$ if $n \equiv 0 \pmod{N}$, and $\delta(n) = 0$ otherwise. Set $[x] = N^{-1} (1 - x^N)/(1 - x)$.

The N -dimensional Faddeev-Kashaev quantum dilogarithm and its inverse are the N^2 -matrices whose components are the rational functions defined on the curve Γ by

$$R(x, y, z)_{\alpha, \beta}^{\gamma, \delta} = h(z/x) \zeta^{\alpha\delta + \frac{\alpha^2}{2}} \omega(x, y, z|\gamma - \alpha) \delta(\gamma + \delta - \beta)$$

$$\bar{R}(x, y, z)_{\gamma, \delta}^{\alpha, \beta} = \frac{[x/z]}{h(z/x)} \zeta^{-\alpha\delta - \frac{\alpha^2}{2}} \frac{\delta(\gamma + \delta - \beta)}{\omega(\frac{x}{z}, y, z|\gamma - \alpha)} \quad .$$

We can interpret these matrices as functions of $\mathcal{I}$ -tetrahedra as follows. Let (Δ, b, w) be an $\mathcal{I}$ -tetrahedron. Write $w_i = -p_{i+1}/p_{i+2}$ (indices mod $(\mathbb{Z}/3\mathbb{Z})$) as after Def. 2.3. Recall that $p_0 + p_1 + p_2 = 0$. Fix a common determination of the N -th roots of the p_i 's, which we denote by p'_i . Set

$$\mathfrak{L}(\Delta, b, w) = \begin{cases} R(p'_1, p'_0, -p'_2) & \text{if } * = 1 \\ \bar{R}(p'_1, p'_0, -p'_2) & \text{if } * = -1 \end{cases} \quad ,$$

where $*$ = ± 1 according to the b -orientation of Δ . Note that (13) implies that $\mathfrak{L}(\Delta, b, w)$ only depends on (b, w) , and not on the N -th roots p'_i of the p_i 's, for it is homogeneous in these variables. One realizes that

$\mathfrak{L}(\Delta, b, w)$ does not respect the tetrahedral symmetries.

Let us consider the 1-skeleton of the cell decomposition of Δ dual to the canonical triangulation with 4 vertices. It is made by 4 edges incident to the center of Δ . They are oriented by the orientation complementary to the b -orientations of the dual 2-faces of Δ . Hence there is a couple of arcs incoming into Δ and a couple of outgoing ones. One can associate to both couples a copy of $\mathbb{C}^N \otimes \mathbb{C}^N$ (with the standard basis) and interpret $\mathfrak{L}(\Delta, b, w)$ as a linear operator defined on the incoming couple, with values in the outgoing one.

Let $\mathcal{T}_{\mathcal{I}} = (T, b, w)$ be any $\mathcal{I}$ -triangulation for (W, ρ) . Let us consider the 1-skeleton C of the cell decomposition dual to T , with edges oriented as above. By associating to each (Δ_i, b_i, w_i) the corresponding operator $\mathfrak{L}(\Delta_i, b_i, w_i)$, one gets an *operator network* whose complete contraction gives a scalar $\mathfrak{L}_N(\mathcal{T}_{\mathcal{I}}) \in \mathbb{C}$ (note that there is no edge with free ends in C). This has an explicit expression as a *state sum*, where the *states* are given by the indices of the matrices entries as follows. A state is a function defined on the edges of C , with values in $\{0, \dots, N-1\}$. For every (Δ_i, b_i, w_i) , any state α determines an entry (a $6j$ -symbol) $\mathfrak{L}(\Delta_i, b_i, w_i)_{\alpha}$ of $\mathfrak{L}(\Delta_i, b_i, w_i)$. Set

$$\mathfrak{L}_N(\mathcal{T}_{\mathcal{I}})_{\alpha} = \prod_i \mathfrak{L}(\Delta_i, b_i, w_i)_{\alpha}$$

and

$$\mathfrak{L}_N(\mathcal{T}_{\mathcal{I}}) = \sum_{\alpha} \mathfrak{L}_N(\mathcal{T}_{\mathcal{I}})_{\alpha} \quad . \quad (14)$$

One realizes that

$\mathfrak{L}_N(\mathcal{T}_{\mathcal{I}})$ is invariant only for some peculiar instances of $\mathcal{I}$ -transits (one being the same as for Schaeffer's identity for the Rogers dilogarithm).

These facts justify the following name: $\mathfrak{L}(\Delta, b, w)$ is the N -dimensional *non symmetric quantum dilogarithm*, computed on the given $\mathcal{I}$ -tetrahedron. The special $\mathcal{I}$ -transits which keep $\mathfrak{L}_N(\mathcal{T}_{\mathcal{I}})$ invariant induce the *basic five term (pentagonal) relations* satisfied by the non symmetric quantum dilogarithm. In order to construct invariants for (W, ρ) based on the quantum dilogarithms $\mathfrak{L}$, one has to solve the same kind of difficulties than for defining the dilogarithmic invariant, based on the classical Rogers dilogarithm L .

5.2 Tetrahedral symmetries

Let (Δ, b, w) be as above. An *integral charge* on (Δ, b, w) is a $\mathbb{Z}$ -valued map on the edges of Δ such that $c(e) = c(e')$ for opposite edges e and e' , and $c_0 + c_1 + c_2 = 1$, where $c_i = c(e_i)$. Note that a charge is formally similar to a flattening. The main difference is that the charge does not depend on the moduli w . In fact a charge defines a flattening only if $*_w = -1$. Write $N = 2p + 1$, and for each edge e set $c'(e) = (p + 1) c(e) \bmod(N)$, viewed as a point in $\{0, \dots, N-1\}$.

Definition 5.1 The *symmetrized quantum dilogarithm* is the matrix-valued operator, defined on the set of charged $\mathcal{I}$ -tetrahedra (Δ, b, w, c) , given by

$$\mathfrak{R}(\Delta, b, w, c) = \begin{cases} ((-p'_1/p'_2)^{-c_1} (-p'_2/p'_0)^{c_0})^p R'(w|c) & \text{if } * = 1 \\ ((-p'_1/p'_2)^{-c_1} (-p'_2/p'_0)^{c_0})^p \bar{R}'(w|c) & \text{if } * = -1 \end{cases} , \quad (15)$$

where $*$ = ± 1 according to the b -orientation of Δ , and the matrices $R'(w|c)$ and $\bar{R}'(w|c)$ have the components

$$R'(w|c)_{\alpha,\beta}^{\gamma,\delta} = \zeta^{c'_1(\gamma-\alpha)} R(p'_1, p'_0, -p'_2)_{\alpha,\beta-c'_0}^{\gamma-c'_0,\delta} \quad (16)$$

$$\bar{R}'(w|c)_{\gamma,\delta}^{\alpha,\beta} = \zeta^{c'_1(\gamma-\alpha)} \bar{R}(p'_1, p'_0, -p'_2)_{\gamma+c'_0,\delta}^{\alpha,\beta+c'_0}.$$

Note again that (13) implies that $\mathfrak{R}(\Delta, b, w, c)$ only depend on (b, w, c) , and not on the choice of the N -th roots p'_i of the p_i 's. Here we identify $1/2 \in \mathbb{Z}/N\mathbb{Z}$ with $p+1$.

Remark 5.2 This symmetrization slightly differs from the one adopted in [2] for B -characters, see Remark 5.7 below. This new one has been obtained by formally copying the flattening's contributions in $\exp(R(\Delta, b, w, f))$, postulating that charges and flattenings should essentially play the same role. In fact, it works.

Recall that the permutation group on four elements, which is the symmetry group of a branched abstract tetrahedron, is generated by the transpositions (01), (12) and (23). The following lemma describes the tetrahedral symmetries of $\mathfrak{R}$.

Lemma 5.3 *Let (Δ, b, w, c) be a charged $\mathcal{I}$ -tetrahedron. Changing the orientation of the edge e_0 , e_1 or e'_0 by the transposition (01), (12) or (23) of its vertices gives respectively*

$$\begin{aligned} \mathfrak{R}((01)(\Delta, b, w, c)) &\equiv_{(\pm\zeta^{\mathbb{Z}})} T_1^{-1} \mathfrak{R}(\Delta, b, w, c) T_1 \\ \mathfrak{R}((12)(\Delta, b, w, c)) &\equiv_{(\pm\zeta^{\mathbb{Z}})} S_1^{-1} \mathfrak{R}(\Delta, b, w, c) T_2 \\ \mathfrak{R}((23)(\Delta, b, w, c)) &\equiv_{(\pm\zeta^{\mathbb{Z}})} S_2^{-1} \mathfrak{R}(\Delta, b, w, c) S_2 \quad , \end{aligned}$$

where $\equiv_{(\pm\zeta^{\mathbb{Z}})}$ means equality up to sign and multiplication by N -th roots of unity. Here we write $T_1 = T \otimes 1$, etc..., and T and S are the N -dimensional invertible square matrices with components $T_{m,n} = \nu \zeta^{\frac{m^2}{2}} \delta(m+n)$ and $S_{m,n} = N^{-\frac{1}{2}} \zeta^{mn}$, where $\nu = g(1)/|g(1)|$.

The proof is based on the same computations that gave Prop. 9.6 in the Appendix of [2], adapted to the new symmetrization.

5.3 Complete five term relations

Replacing the non symmetric quantum dilogarithms with the symmetrized ones in (14) for charged $\mathcal{I}$ -triangulations (T, b, w, c) of (W, ρ) , one obtains state sums $\mathfrak{R}_N(\mathcal{T}_T, c)$. The next step is to complete the $\mathcal{I}$ -transits to suitable charged $\mathcal{I}$ -transits, in order to realize the full transit invariance of the state sum's value. As the charges are moduli-independent, also their transit is.

One says that there is a *charge transit* $(T, c) \leftrightarrow (T', c')$ if c' equals c on the edges of the abstract tetrahedra of T not involved in the move, and for any other edge e we have the *transit of sum* condition:

$$\sum_{a \in \epsilon_T^{-1}(e)} c(a) = \sum_{a' \in \epsilon_{T'}^{-1}(e)} c'(a') \quad .$$

Note that, for $2 \rightarrow 3$ transits, this implies that the sum of the charges around the new edge after the move is equal to 2. Note also that a charge transit coincides with a flattening transit, providing that the signs $*_b$'s and $*_w$'s satisfy certain conditions which are easy to determine. By using Lemma 5.3 and the basic five term (pentagonal) relations for the non symmetric quantum dilogarithms we finally get:

Proposition 5.4 *For any charged $2 \leftrightarrow 3$ $\mathcal{I}$ -transit $(T, w, c) \leftrightarrow (T', w', c')$ we have*

$$\prod_{\Delta_i \subset T} \mathfrak{R}(\Delta_i, b, w, c) \equiv_{(\pm \zeta^z)} \prod_{\Delta'_i \subset T'} \mathfrak{R}(\Delta'_i, b', w', c') \quad .$$

5.4 Link-fixing and QHI for triples (W, L, ρ)

As for flattenings in the definition of the dilogarithmic invariant, the next step is to give the right notion of global integral charge which should be stable for charged $\mathcal{I}$ -transits. The solution of this problem is much more elaborated than for the dilogarithmic invariant. The naive idea should be to require that the sum of charges around each edge of T is equal to 2. But simple combinatorial considerations show that such tentative global integral charges do not exist. A way to overcome this difficulty is to fix an arbitrary link L in W (considered up to ambient isotopy) and to inglobe it in all the construction of the QHI.

A $\mathcal{D}$ -triangulation $(\mathcal{T}, H)$ for (W, L, ρ) is a $\mathcal{D}$ -triangulation $\mathcal{T}$ for (W, ρ) with a Hamiltonian (i.e. which contains all the vertices of T) subcomplex H which realizes the link L . An integral charge c on $(\mathcal{T}, H)$ satisfies by definition the following conditions:

- (1) the sum of its values equal 2 (resp. 0) around the edges of $T \setminus H$ (resp. H);
- (2) it satisfies the same $\mathbb{Z}/2\mathbb{Z}$ -cohomological condition than for flattenings: there is a class $[c] \in H^1(W; \mathbb{Z}/2\mathbb{Z})$, and we require that $[c] = 0$.

Note that any charge c eventually encodes H , hence the link L . The existence of charged $\mathcal{D}$ - and $\mathcal{I}$ -triangulations $(\mathcal{T}, c)$, $(\mathcal{T}_{\mathcal{I}}, c)$ for (W, L, ρ) is a rather demanding fact proved in [1] and [2]. Again, the good affine structure on the set of charges is a variation of the fundamental Neumann's result Th. 4.2. The presence of the link must be integrated also in the transits; in particular the condition to be Hamiltonian must be preserved. For instance, for $2 \rightarrow 3$ transits the new edge must be in $T' \setminus H'$; for positive bubble moves one edge of the initial triangle must belong to H , and it is replaced with the two edges connected to it and to the new vertex in T' . Finally one achieves the construction of the QHI:

Theorem 5.5 *The value of the (normalized) state sum*

$$H_N(\mathcal{T}_{\mathcal{I}}, c) = N^{-n_0} \mathfrak{R}_N(\mathcal{T}_{\mathcal{I}}, c) \quad ,$$

where n_0 is the number of vertices of T , does not depend on the choice of the charged $\mathcal{I}$ -triangulation $(\mathcal{T}_{\mathcal{I}}, c)$ for (W, L, ρ) , up to sign and multiplication by N -th roots of unity. Hence, up to this ambiguity, for any triple (W, L, ρ) it defines a quantum hyperbolic invariant $H_N(W, L, \rho) \in \mathbb{C}$.

Remark 5.6 The presence of the link L and the ambiguity up to multiplication by N -th roots of unity of $H_N(W, L, \rho)$ are entirely a consequence of the specific symmetrization procedure of the quantum dilogarithms we have adopted. We guess that suitable variations of this procedure (by using moduli-dependent charges, or by following even more closely the strategy used for uniformizing the Rogers dilogarithm) allows one to define the QHI directly for (W, ρ) , and also to well define them only up to multiplication by $\exp(i\pi/12)$. We postpone this problem to a future work.

Remark 5.7 The QHI for B -characters defined in [2] and their state sum formulas differ from those in 5.5 by a scalar factor depending on the cocycle z of $\mathcal{T}$, not only on $\mathcal{T}_{\mathcal{I}}$. This is a consequence of the different symmetrization adopted in [2]. There, it consisted in replacing in (15) the scalar state-independent factor in front of the

matrices R' and $\bar{R}'$ by $(-q'_2)^p$, where the q_j 's have been defined in Remark 2.5 (3) and $'$ denotes, as before, a common determination of the N -th roots of the q_j 's. Let us denote by $\mathfrak{R}^B(\mathcal{T}, c)$ the associated state sum.

Then, the statement of Lemma 5.3 is unchanged, except that there is no sign ambiguity. However, in Prop. 5.4 one has to multiply both sides by the respective $Q_2 := \prod_i (-q'_2)_i^p$. It is a remarkable but somewhat fortuitous fact that, for B -characters and for any positive $2 \rightarrow 3$ $\mathcal{D}$ -transit $\mathcal{T} \rightarrow \mathcal{T}'$, one has $Q_2(\mathcal{T}')/Q_2(\mathcal{T}) = x(e)^{2p}$, where $x(e)$ is the upper-diagonal value of the cocycle z on the new edge in $\mathcal{T}' \setminus H'$. Normalizing $\mathfrak{R}^B(\mathcal{T}, c)$ by dividing it by $\prod_{e \in \mathcal{T} \setminus H} x(e)^{2p}$, it eventually gives a well defined invariant up to N -th roots of unity. The same procedure for general $PSL(2, \mathbb{C})$ -characters (using the p'_2 's instead of the q'_2 's) does not seem to work, due to the fact that the explicit formula for $P_2(\mathcal{T}')/P_2(\mathcal{T})$ heavily depends on the branching.

6 Scissors congruence classes

Consider the free $\mathbb{Z}$ -modules $\mathbb{Z}(\mathcal{D})$ and $\mathbb{Z}(\mathcal{I})$ which are respectively generated by all $\mathcal{D}$ -tetrahedra and all $\mathcal{I}$ -tetrahedra. Let $\mathcal{P}(\mathcal{D})$ (resp. $\mathcal{P}(\mathcal{I})$) be the quotient of $\mathbb{Z}(\mathcal{D})$ (resp. $\mathbb{Z}(\mathcal{I})$) by all instances of the relations associated to the tetrahedral symmetries in Lemma 2.7, and to the $2 \leftrightarrow 3$ $\mathcal{D}$ -transits (resp. $\mathcal{I}$ -transits).

We call $\mathcal{P}(\mathcal{D})$ (resp. $\mathcal{P}(\mathcal{I})$) the $\mathcal{D}$ - (resp. $\mathcal{I}$ -) *(pre)-Bloch group*. Prop. 2.14 implies that there is a surjective homomorphism $I : \mathcal{P}(\mathcal{D}) \rightarrow \mathcal{P}(\mathcal{I})$.

Working with either flattened or charged $\mathcal{I}$ -tetrahedra, and using the appropriate lifts of the tetrahedral symmetries and of the five term identities, one obtains the F - and C -refined $\mathcal{I}$ -(pre)-Bloch groups $\mathcal{P}(\mathcal{I})_F$ and $\mathcal{P}(\mathcal{I})_C$.

Clearly, every $\mathcal{D}$ -triangulation $\mathcal{T}$ and every $\mathcal{I}$ -triangulation $\mathcal{T}_{\mathcal{I}}$ for (W, ρ) defines an element $\mathfrak{c}(\mathcal{T}) \in \mathcal{P}(\mathcal{D})$ and $\mathfrak{c}(\mathcal{T}_{\mathcal{I}}) \in \mathcal{P}(\mathcal{I})$ respectively. Similarly, any flattened $(\mathcal{T}_{\mathcal{I}}, f)$ defines an element $\mathfrak{c}(\mathcal{T}_{\mathcal{I}}, f) \in \mathcal{P}(\mathcal{I})_F$, and any charged $(\mathcal{T}_{\mathcal{I}}, c)$ for (W, L, ρ) defines an element $\mathfrak{c}(\mathcal{T}_{\mathcal{I}}, c) \in \mathcal{P}(\mathcal{I})_C$. One has

Proposition 6.1 *The elements $\mathfrak{c}(\mathcal{T})$, $\mathfrak{c}(\mathcal{T}_{\mathcal{I}})$, $\mathfrak{c}(\mathcal{T}_{\mathcal{I}}, f)$ and $\mathfrak{c}(\mathcal{T}_{\mathcal{I}}, c)$ do not depend on the choices in their respective arguments. Hence they define invariants $\mathfrak{c}_D(W, \rho) \in \mathcal{P}(\mathcal{D})$, $\mathfrak{c}_I(W, \rho) \in \mathcal{P}(\mathcal{I})$, $\mathfrak{c}_F(W, \rho) \in \mathcal{P}(\mathcal{I})_F$ and $\mathfrak{c}_C(W, L, \rho) \in \mathcal{P}(\mathcal{I})_C$.*

These are generally called scissors congruence classes of (W, ρ) or of (W, L, ρ) . Moreover, the homomorphism I sends $\mathfrak{c}_D(W, \rho)$ onto $\mathfrak{c}_I(W, \rho)$.

Clearly, the formula in Th. 4.3 defines a function $R : \mathcal{P}(\mathcal{I})_F \rightarrow \mathbb{C}/(\pi^2/6)\mathbb{Z}$ such that

$$R(W, \rho) = R(\mathfrak{c}_F(W, \rho)) .$$

One would like to interpret also the QHI as evaluations on $\mathfrak{c}_C(W, L, \rho)$ of suitable functions defined on $\mathcal{P}(\mathcal{I})_C$. This is roughly true, but there is some subtilities (see the discussion at the end of Section 5 of [4]).

Lemma 2.10 implies that every $\mathcal{I}$ -triangulation $\mathcal{T}_{\mathcal{I}}$ for (W, ρ) also defines a class $\alpha(\mathcal{T}_{\mathcal{I}}) \in H_3(PSL(2, \mathbb{C}), \mathbb{Z})$, where $PSL(2, \mathbb{C})$ is endowed with the discrete topology. One can prove that also $\alpha(\mathcal{T}_{\mathcal{I}})$ does not depend on the choice of $\mathcal{T}_{\mathcal{I}}$, hence it defines an invariant $\alpha(W, \rho) \in H_3(PSL(2, \mathbb{C}), \mathbb{Z})$. This is a main ingredient of the group-cohomological approach of Dupont-Sah.

7 Further comments on the proofs

The proofs of Th. 4.3 and Th. 5.5 have the very same structure. In fact one follows, almost *verbatim*, the proof given in {Sections 1-6 + the Appendix} of [2] for the

B-QHI. For the dilogarithmic invariant the proof is easier, because one does not have to keep track of the link L during the transits.

A key point is that quasi-regular triangulations can be connected by quasi-regular transits. This is proved in Prop. 2.10 of [2]. This implies that also $\mathcal{D}$ -transits and $\mathcal{I}$ -transits are *generically* possible, and this is enough. In fact, another delicate point is the invariance w.r.t. the flattenings or the charges. This heavily depends on Neumann's crucial Th. 4.2 and runs like in the proof of Th. 4.6 in [2].

For scissors congruence classes, the above *generic* existence of $\mathcal{D}$ -transits and $\mathcal{I}$ -transits is not enough. We have to refine Prop. 2.10 of [2] (by using the same technics indeed), in order to get the full existence of transits. There is another subtlety: the (pre)-Bloch groups are defined by only using the relations coming from $2 \leftrightarrow 3$ transits. But to connect triangulations (with possibly a different number of vertices), one has to use also the 'bubble' transits. So one has to check that this induces relations which are in fact consequences of the other ones. This is proved as in Corollary 4.5 of [2].

The situation is a little more complicated if one works with arbitrary branchings, or if one allows also *degenerate* ideal tetrahedra. General branchings were already treated in [2], so one simply repeats those arguments. In presence of degenerate tetrahedra, the invariance of the dilogarithmic sum w.r.t. the branching becomes more delicate. This is automatic under the non-degenerate assumption, thanks to Lemma 3.3. Anyway, one can obtain it by using the quasi-regular refinement of the following result of F. Costantino [7]:

Two branchings on a same given triangulation T of W can be connected by transits of branchings.

This allows to reduce also the branching invariance to the invariance by transits.

8 A Volume Conjecture for the QHI of (W, L, ρ)

It is clear from the discussion in Section 5 that the structure of the QHI is modeled on the one of $\exp((1/2i\pi)\mathcal{R}(W, \rho))$. In fact one introduces the $1/2i\pi$ -factor to have a perfect agreement of the behaviour with respect to the involution $W \rightarrow -W$, given by the change of orientation of W :

Lemma 8.1 *Denote by * is the complex conjugation. We have*

$$\begin{aligned} H_N(-W, L, \rho) &= (H_N(W, L, \rho^*))^* \\ \exp((1/2i\pi)\mathcal{R}(-W, \rho)) &= (\exp((1/2i\pi)\mathcal{R}(W, \rho^*)))^* \end{aligned}$$

The first identity is proved like in Prop. 6.1. in [2]. The second one is a consequence of Prop. 4.4, and of $\text{CS}(\rho^*) = \text{CS}(\rho)$ and $\text{Vol}(\rho^*) = -\text{Vol}(\rho)$. These structural coincidences and the actual asymptotic behaviour of the quantum dilogarithms (see below) motivate the following *Volume Conjecture* for the asymptotic expansion of QHI, when $N \rightarrow \infty$:

Conjecture 8.2 *There exist invariants $C = C(W, L, \rho) \in \mathbb{C} \bmod(\pi^2/6)\mathbb{Z}$ and $D = D(W, L, \rho) \in \mathbb{C}^*$ such that*

$$(H_N(W, L, \rho))^N \equiv \left[\exp\left(\frac{C + N\mathcal{R}(W, \rho)}{2i\pi}\right) \right]^N \left(D + \mathcal{O}\left(\frac{1}{N}\right) \right)$$

where $\equiv$ means equality up to multiplication by the integer powers of $\exp(i\pi/12)$.

Conjecture 8.2 says at first that $H_N(W, L, \rho)^N$ has an exponential growth rate. Assuming it, the fact that $\exp(C/i\pi)$, $\exp(R(W, \rho)/i\pi)$ and D are well-determined invariants of (W, L, ρ) follows from the invariance of $H_N(W, L, \rho)^N$ and the uniqueness of the coefficients of asymptotic power series expansions.

At present, the nature of C and D is somewhat mysterious to us. There are no reasons to expect that, for instance, $C = 0$ or $D = 1$. We have expressed the conjecture in terms of the N -th power of $H_N(W, L, \rho)$ so as to kill its multiplicative ambiguity up to N -th roots of unity. (The statement is formally the same as in [4], except that it was given mod $(i\pi/4)\mathbb{Z}$, see the discussion at the end of Section 4).

Classical manipulations of one-variable complex analysis with the so-called Faddeev's *non-compact dilogarithm* [11] allows one to prove that when $N \rightarrow \infty$ (we use the functions introduced in section 5.1):

$$g(z/x) \omega(x, y, z|n) \sim (y/z)^n \exp \left[\frac{N}{2i\pi} (\text{Li}_2((x/z)\zeta^n)) + \log(x/z)^2 - \pi \log(x/z) + \pi^2 \right],$$

where $\log$ is, as before, the standard branch of the logarithm. Rewriting n in terms of states and charges, one derives from this formula the leading term of the asymptotic expansion of the symmetrized quantum dilogarithms. This corroborates Conjecture 8.2.

Remarks 8.3 The above Volume Conjecture predicts, in particular, that the dominant term of the asymptotic expansion of the QHI for $N \rightarrow \infty$ is *not*-sensitive to the link-fixing used to get one specific global symmetrization of the state sums. This is not clear for C and D . There is a simple way to modify the dilogarithmic invariant so as to make it link-sensitive. For that, it is enough to set

$$R(W, L, \rho) = R(W, \rho) + (i\pi/2) \log(\text{Tr}(\alpha(L))) \quad ,$$

where α is any representative of ρ and L is arbitrarily oriented, and considered as an element of the fundamental group of W . Note that $R(W, L, \rho)$ does not depend on the choices we made. It can be computed by using any $\mathcal{D}$ -triangulation $\mathcal{T}$ for (W, L, ρ) , but not only by using the idealization $\mathcal{T}_{\mathcal{I}}$.

9 Cusped manifolds

This section is less definitive than the rest of the paper. The final achievement of the results presented below is strictly related to the solution of the problem mentioned in Remark 5.6.

9.1 Dilogarithmic invariant and QHI

Let M be an oriented complete non-compact and finite volume hyperbolic 3-manifold (shortly: M is a *cusped manifold*). Fix a triangulation of M made by embedded geodesic ideal tetrahedra. For simplicity, assume that it is branched. This is not really necessary here, due to the symmetry relations in Lemma 3.3 and Lemma 5.3; on the contrary, for the case of (W, ρ) treated before, the branching was part of the structure of $\mathcal{D}$ -triangulations, which dominate the $\mathcal{I}$ -triangulations.

With the usual notations, this triangulation of M can be represented as $\mathcal{T}_{\mathcal{I}} = (T, b, w)$ such that every (Δ_i, b_i, w_i) is *quasi-geometric*, i.e. geometric in the sense of Def. 2.9 or possibly degenerate.

One can enrich $\mathcal{T}_{\mathcal{I}}$ with suitable flattenings f or integral charges c (again this is essentially due to [18]). Their definition is similar to the one given for (W, ρ) , but

there is a further condition which mimics the usual *completeness condition* satisfied by the moduli. So one can define both $R(\mathcal{T}_I, f) \in \mathbb{C}/(\pi^2/6)\mathbb{Z}$ and $H_N(\mathcal{T}_I, c)$.

One would like to prove that they do not depend on any choice, hence that they define invariants for M , just by using our direct methods based on $\mathcal{I}$ -transits. It is not clear to us if any two such enriched quasi-geometric $\mathcal{I}$ -triangulations of M can be connected by enriched $\mathcal{I}$ -transits. However, for $R(\mathcal{T}_I, f)$ one can prove it if we impose for instance the following further restriction on $\mathcal{T}_I$:

Proposition 9.1 *Suppose that M admits triangulations (T, b) such that the volume function, defined on the set of all $\mathcal{I}$ -triangulations supported by (T, b) , has a unique maximum at $\mathcal{T}_I$. Then $R(\mathcal{T}_I, f)$ does not depend on any choice, thus defining a dilogarithmic invariant $R(M) \in \mathbb{C} \bmod (\pi^2/6)\mathbb{Z}$.*

Here, the set of all $\mathcal{I}$ -triangulations supported by (T, b) includes *all* (non necessarily quasi-geometric) solutions of the compatibility and completeness equations supported by T . It is not known to us if any cusped M has such special triangulations, but, as is used in the Snappea program it often happens.

The same result should hold true for the expected link-free definition of QHI (see Remark 5.6); then, let us assume that also the QHI $H_N(M)$ of M are well-defined. The Volume Conjecture 8.2 for (W, ρ) can be repeated *verbatim* for these invariants of cusped manifolds.

9.2 About the Volume Conjecture for the colored Jones invariants $J_N(L)$

Recall that Kashaev's Volume Conjecture [15], reformulated in terms of the (suitably normalized) colored Jones invariants $J_N(L)$ of links in S^3 by Murakami-Murakami, states that for an hyperbolic knot L we have

$$\lim_{N \rightarrow \infty} (2\pi/N) \log(|J_N(L)|) = \text{Vol}(M) \quad ,$$

where M is the cusped manifold given by the hyperbolic complement of L in S^3 . Due to Remark 8.3, one has:

The Volume Conjecture 8.2 for $H_N(W, L, \rho)$ is not compatible with the one stated above for $J_N(L)$ if one also assumes (as it is currently done, see [14]) that $J_N(L)^N$ coincides with $H_N(S^3, L, \rho_0)^N$, where ρ_0 is the necessarily trivial character.

We guess that:

- (1) the last assumption above is not correct (we will elaborate on this point in [3]).
- (2) The Volume Conjecture for $J_N(L)$ makes sense only if it could be related to Conjecture 8.2 for general cusped manifolds. At least they are formally compatible. However, for the moment, we do not see a systematic way to identify (even asymptotically) $J_N(L)$ and $H_N(M)$, where M is the cusped manifold $S^3 \setminus L$.
- (3) A consistent relationship between all these Volume Conjectures could be obtained thanks to Thurston's *hyperbolic Dehn filling* and *double limit procedure*. Indeed, let M be a cusped manifold and (W_n, L_n, ρ_n) be a sequence of compact hyperbolic Dehn fillings of M converging to M . Here, L_n denotes the link made of the short simple geodesics in W_n forming the *cores* of the fillings, and ρ_n is the holonomy of the hyperbolic manifold W_n . Presumably, one has

$$R(W_n, \rho_n) \longrightarrow R(M) \quad , \quad n \rightarrow \infty \quad ,$$

and for every fixed N also

$$H_N(W_n, L_n, \rho_n) \longrightarrow H_N(M) \quad , \quad n \rightarrow \infty \quad .$$

So, following Conjecture 8.2, by taking a 'double limit' we are led to:

Conjecture 9.2 *The dominant term of the asymptotic expansion when $n, N \rightarrow \infty$ of $(H_N(W_n, L_n, \rho_n))^N$ is equal to $\exp((N^2/2i\pi)R(M))$, up to multiplication by integer powers of $\exp(i\pi/12)$.*

The Volume Conjecture for J_N seems to be corroborated by few numerical computations, obtained via a quite formal and ‘optimistic’ use of the *stationary phase method*. We also guess that the above considerations could help to have a correct understanding of the *meaning* of those formal manipulations.

References

- [1] S. Baseilhac, *Dilogarithme Quantique et Invariants de 3-Variétés*, Thèse, Université Paul Sabatier, Toulouse (France), Octobre 2001;
- [2] S. Baseilhac, R. Benedetti, *QHI Theory, I: 3-manifold scissors congruence classes and quantum hyperbolic invariants*, arXiv.math.GT/0201240;
- [3] S. Baseilhac, R. Benedetti, *QHI Theory III: R-Matrices state sums for links in S^3 and the coloured Jones polynomials*, in preparation;
- [4] S. Baseilhac, R. Benedetti, *QHI, 3-manifolds scissors classes and the volume conjecture*, Geometry and Topology Monographs, Volume 4: Invariants of Knots and 3-manifolds (Kyoto 2001), pages 13-28;
- [5] R. Benedetti and C. Petronio, *Lectures on Hyperbolic Geometry*, Springer (1992), 1st edition;
- [6] R. Benedetti and C. Petronio, *Branched Standard Spines of 3-manifolds*, Lect. Notes in Math. No 1653, Springer (1997);
- [7] F. Costantino, PhD Thesis, Scuola Normale Superiore, Pisa;
- [8] J.L. Dupont, C-H. Sah. *Scissors congruences II*, J. Pure and App. Algebra 44 (1987), 137-164;
- [9] J.L. Dupont, *The dilogarithm as a characteristic class for flat bundles*, J. Pure and App. Algebra 44 (1987), 137-164;
- [10] J.L. Dupont, *Scissors Congruences, Group Homology and Characteristic Classes*, Nankai Tracts in Math. Vol.1, World Scientific (2001).
- [11] L.D. Faddeev, *Discrete Heisenberg-Weyl group and modular group*, Lett. Math. Phys. 34 (1995), 249-254;
- [12] L.D. Faddeev, R.M. Kashaev. *Quantum Dilogarithm*, Mod. Phys. Lett. A Vol. 9, No 5 (1994), 427-434;
- [13] R.M. Kashaev. *Quantum dilogarithm as a 6j-symbol*, Mod. Phys. Lett. A Vol. 9, No 40 (1994), 3757-3768;
- [14] R.M. Kashaev, *A link invariant from quantum dilogarithm*, Mod. Phys. Lett. A Vol. 10 (1995), 1409-1418;
- [15] R.M. Kashaev, *The hyperbolic volume of knots from the quantum dilogarithm*, Lett. Math. Phys. 39 (1997), 269-275;
- [16] L.Lewin, *Dilogarithm and Associated Functions*, Mc Donald (1958);

- [17] S.V. Matveev, *Transformations of special spines and the Zeeman conjecture*, Math. USSR Izvestia No 31 (1988), 423-434;
- [18] W.D. Neumann, *Combinatorics of triangulations and the Chern-Simons invariant for hyperbolic 3-manifolds*, in *Topology'90* (Columbus, OH, 1990), De Gruyter, Berlin (1992);
- [19] W.D. Neumann, *Hilbert's 3rd problem and invariants of 3-manifolds*, The Epstein birthday schrift, Geom. Topol. Monographs Vol. 1, Coventry (1998), 383-411;
- [20] W.D. Neumann, *Extended Bloch group and the Chern-Simons class*, incomplete working version, <http://www.math.columbia.edu/~neumann>;
- [21] W.D. Neumann, J. Yang. *Bloch invariants of hyperbolic 3-manifolds*, Duke Math. Journal Vol. 96, No 1 (1999);
- [22] R. Piergallini, *Standard moves for standard polyhedra and spines*, Rend. Circ. Mat. Palermo No 37, suppl. 18 (1988), 391-414;
- [23] L.J. Rogers, *On function sums connected with the series $\sum x^n/n^2$* , Proc. London Math. Soc. 4 (2) (1907), 169-189;
- [24] D. Zagier, *The remarkable dilogarithm*, Jour. Math. Phys. Sci., Vol. 22, No 1 (1988), 131-145;